



## SOLVABLE AND/OR INTEGRABLE MANY-BODY MODELS ON A CIRCLE\*

OKSANA BIHUN and FRANCESCO CALOGERO<sup>†</sup>

*Department of Mathematics, Concordia College, 901 8th Str. South  
Moorhead, MN, USA*

<sup>†</sup> *Physics Department, University of Rome “La Sapienza”, Rome, Italy*

**Abstract.** Various many-body models are treated, which describe  $N$  points confined to move on a plane circle. Their Newtonian equations of motion (accelerations equal forces) are *integrable*, i.e., they allow the *explicit* exhibition of  $N$  *constants of motion* in terms of the dependent variables and their time-derivatives. Some of these models are moreover *solvable* by purely algebraic operations, by (explicitly performable) quadratures and, finally, by functional inversions. The techniques to manufacture these models are not new, but some of these models are themselves new and others are reinterpretations of known models.

### 1. Introduction

The investigation of the time evolution of an arbitrary number  $N$  of point-particles the dynamics of which is determined by Newtonian equations of motion (accelerations equal forces) is of course a fundamental topic in physics and mathematics. The identification in this context of models *amenable to exact treatments* is a major area of research in mathematical physics and applied mathematics, having a centuries-old history and having been boosted by developments in the last few decades, which also impacted several areas of physics beyond mechanics and many fields of pure mathematics. An interesting related development which is now becoming of interest is the study of such models in which the motion is restricted to lie on an *a priori* prescribed manifold: see for instance [1, 5, 6, 8]. In this paper we make some initial, simple steps in this direction by focussing on

---

\*Reprinted from J. Geom. Symmetry Phys. **30** (2013) 1–18.